Analysis: Women's pain is routinely underestimated, and gender stereotypes are to blame

9 April 2021

The suspicion that gender stereotypes could lead doctors to underestimate women's pain has been confirmed by research which found healthcare staff, both men and women, often discount women's pain, says Professor Amanda Williams (UCL Clinical, Education & Health Psychology).





Editorial | Published: 11 October 2024

Funding research on women's health

Nature Reviews Bioengineering 2, 797–798 (2024) Cite this article

9822 Accesses 2 Citations 43 Altmetric Metrics

Women's health research has long been underfunded, in part owing to stigmas associated with conditions that primarily affect women. Equitable health research funding requires transparency from funding agencies, investment in women-centred innovations, support for women in science and a cultural shift in how health issues are viewed.

Research related to women's health is critically underfunded. In 2020, only 5% of global research and development (R&D) funding was allocated to women's health research. This was ARTS & ENTERTAINMENT, HEALTH, NATIONAL

ChatGPT Thinks Doctors Are Male—How Al Mirrors Society's **Gender Bias**

PUBLISHED 8/1/2024 by SHIKHA JAIN and VINEET ARORA



Gender bias revealed in AI tools screening for liver disease

11 July 2022

Artificial intelligence models built to predict liver disease from blood tests are twice as likely to miss disease in women as in men, a new study by UCL researchers has found.









